

Exact Chromatic Polynomials for Toroidal Chains of Complete Graphs

Shu-Chiuan Chang*

C. N. Yang Institute for Theoretical Physics
State University of New York
Stony Brook, N. Y. 11794-3840

Abstract

We present exact calculations of the partition function of the zero-temperature Potts antiferromagnet (equivalently, the chromatic polynomial) for graphs of arbitrarily great length composed of repeated complete subgraphs K_b with $b = 5, 6$ which have periodic or twisted periodic boundary condition in the longitudinal direction. In the $L_x \rightarrow \infty$ limit, the continuous accumulation set of the chromatic zeros $\mathcal{B}$ is determined. We give some results for arbitrary b including the extrema of the eigenvalues with coefficients of degree $b - 1$ and the explicit forms of some classes of eigenvalues. We prove that the maximal point where $\mathcal{B}$ crosses the real axis, q_c , satisfies the inequality $q_c \leq b$ for $2 \leq b$, the minimum value of q at which $\mathcal{B}$ crosses the real q axis is $q = 0$, and we make a conjecture concerning the structure of the chromatic polynomial for Klein bottle strips.

*email: shu-chiuan.chang@sunysb.edu

I. INTRODUCTION

The q -state Potts antiferromagnet (AF) [1,2] exhibits nonzero ground state entropy, $S_0 > 0$ (without frustration) for sufficiently large q on a given lattice Λ or, more generally, on a graph G . This is equivalent to a ground state degeneracy per site $W > 1$, since $S_0 = k_B \ln W$. There is a close connection with graph theory here, since the zero-temperature partition function of the above-mentioned q -state Potts antiferromagnet on a graph G satisfies

$$Z(G, q, T = 0)_{PAF} = P(G, q) , \quad (1.1)$$

where $P(G, q)$ is the chromatic polynomial expressing the number of ways of coloring the vertices of the graph G with q colors such that no two adjacent vertices have the same color (for reviews, see [3]- [5]). The minimum number of colors necessary for such a coloring of G is called the chromatic number, $\chi(G)$. Thus

$$W(\{G\}, q) = \lim_{n \rightarrow \infty} P(G, q)^{1/n} , \quad (1.2)$$

where n is the number of vertices of G and $\{G\} = \lim_{n \rightarrow \infty} G$. Where no confusion will result, we shall sometimes write G rather than $\{G\}$ for the infinite-length limit of a given type of strip graph. At certain special points q_s (typically $q_s = 0, 1, \dots, \chi(G)$), one has the noncommutativity of limits

$$\lim_{q \rightarrow q_s} \lim_{n \rightarrow \infty} P(G, q)^{1/n} \neq \lim_{n \rightarrow \infty} \lim_{q \rightarrow q_s} P(G, q)^{1/n} , \quad (1.3)$$

and hence it is necessary to specify the order of the limits in the definition of $W(\{G\}, q_s)$ [6]. Denoting W_{qn} and W_{nq} as the functions defined by the different order of limits on the left and right-hand sides of (1.3), we take $W \equiv W_{qn}$ here; this has the advantage of removing certain isolated discontinuities that are present in W_{nq} .

Using the expression for $P(G, q)$, one can generalize q from $\mathbb{Z}_+$ to $\mathbb{C}$. The zeros of $P(G, q)$ in the complex q plane are called chromatic zeros; a subset of these may form an accumulation set in the $n \rightarrow \infty$ limit, denoted $\mathcal{B}$, which is the continuous locus of points where $W(\{G\}, q)$ is nonanalytic.¹ The maximal region in the complex q plane to which one can analytically continue the function $W(\{G\}, q)$ from physical values where there is nonzero ground state entropy is denoted R_1 . The maximal value of q where $\mathcal{B}$ intersects the (positive) real axis is labeled $q_c(\{G\})$. This point is important since $W(\{G\}, q)$ is a real analytic function from large values of q down to $q_c(\{G\})$.

¹For some families of graphs $\mathcal{B}$ may be null, and W may also be nonanalytic at certain discrete points.

Here we present exact calculations of the partition function of the chromatic polynomial $P(G, q)$ for graphs of arbitrarily great length composed of repeated complete subgraphs K_b with $b = 5, 6$ which have periodic or twisted periodic boundary conditions in the longitudinal direction.² Thus, consider L_x copies of the complete graph K_{L_y} with vertex set $V(K_{L_y}) = \{1, 2, \dots, L_y\}$. Denote the edges joining each adjacent pair of K_{L_y} and K'_{L_y} graphs as $L = \{v_i v'_j | i, j = 1, \dots, L_y\}$, where v_i, v'_j are vertices of adjacent complete graphs K_{L_y} and K'_{L_y} , and impose periodic boundary conditions in the longitudinal direction, x . In [7], the graph with L_x copies of K_{L_y} for arbitrary L_x and L_y was called the bracelet strip. The total number of vertices is $n = L_x L_y$ for these strips.

A generic form for chromatic polynomials for recursively defined families of graphs (families of graphs that can be constructed via repeated addition of some subgraph), of which strip graphs G_s are special cases, is [8]

$$P((G_s)_m, q) = \sum_{j=1}^{N_{G_s, \lambda}} c_{G_s, j}(q) (\lambda_{G_s, j}(q))^{L_x}, \quad (1.4)$$

where $c_{G_s, j}(q)$ and the $N_{G_s, \lambda}$ terms $\lambda_{G_s, j}(q)$ depend on the type of strip graph G_s but are independent of L_x .

For a given type of strip graph G_s , we denote the sum of the coefficients $c_{G_s, j}$ as

$$C(G_s, q) = \sum_{j=1}^{N_{G_s, \lambda}} c_{G_s, j}(q). \quad (1.5)$$

Some works on calculations of chromatic polynomials for recursive families of graphs include [9]- [49].

II. FAMILY OF TOROIDAL STRIPS

Specifically, we consider the set of edges linking two successive complete graphs on the chain is $L = \{11, 22, 33, \dots, L_y L_y\}$, which will be denoted as a torus. For $L_y = 2$, this is the ladder graph with vertex set $T_2 = K_2$ [9]. For the graph with $L_y = 3, 4$, which has vertex set $C_3 = K_3$ and K_4 , the chromatic polynomial was given in [10] and [11], respectively. In [58, 59], these families of graphs with linking edge sets $\{11, 22, 33, \dots, bb\}$ were denoted as

² K_n denotes the complete graph, i.e. the graph with n vertices such that each vertex is connected by an edge to all of the other vertices.

$B_m(b)$, where $b = L_y$, $m = L_x$. The corresponding Klein bottle strip is the same as the toroidal strip except one of the linking edge set is $\{1L_y, 2(L_y - 1), \dots, L_y 1\}$. The form of the chromatic polynomials for the $B_m(b)$ strips was determined to be

$$P(B_m(b), q) = \sum_{d=0}^b \sum_{j=1}^{N_{B_m(b),d,\lambda}} c_{B_m(b),d,j}(q) c'_{B_m(b),d,j} (\lambda_{B_m(b),d,j}(q))^m, \quad (2.1)$$

where d is the degree of the coefficient $c_{B_m(b),d,j}(q)$ as a polynomial in q , and the $c'_{B_m(b),d,j}$'s are numbers which depend on b . The structural feature that these coefficients can be grouped into sets of fixed degree d is similar to that found in [14] for cyclic strips of the square and triangular lattice, although for these bracelet chain graphs, for $b \geq 3$, there is more than one coefficient of a given degree (see [10] for the case $b = 3$), whereas in contrast, for the cyclic strip graphs studied in [14] (and the self-dual family studied in [55]) there is only one coefficient of each degree. The degree d is denoted as the ‘level’ in [60,58,59]. The $\lambda_{B_m(b),d,j}(q)$ is the eigenvalue of an appropriately defined transfer matrix [37] and is a polynomial in q of degree $b - d$. For sufficiently large integer q , the coefficient $c_{B_m(b),d,j}(q) c'_{B_m(b),d,j}$ can be interpreted as the multiplicity of this eigenvalue.

There are partitions associated with each d which were used to determine the forms of $c_{B_m(b),d,j}$ in [59]. For a partition X of d such that $x_d + x_{d-1} + \dots + x_1 = d$ and $x_d \leq x_{d-1} \leq \dots \leq x_1$ where x_i for $2 \leq i \leq d$ are non-negative integers and x_1 is a positive integer, there is an associated partition Y of $d(d+1)/2$ such that $y_d + y_{d-1} + \dots + y_1 = d(d+1)/2$ and $y_i = x_i + d - i$ for $1 \leq i \leq d$. Then the $c_{B_m(b),d,j}$ for a partition X can be written as

$$c_{X,d} = \frac{1}{d!} \prod_{i=1}^d (q - y_i). \quad (2.2)$$

For the partitions $[d]$ and $[1^d]$, the associated $\lambda_{B_m(b),d,j}(q)$'s were given in [58].

A. $L_y = 5$ toroidal strip

By using theorems in [58], we find the chromatic polynomial for $B_m(5)$, i.e., the $5 \times L_x$ strips with each transverse slice forming a K_5 graph, and with edge linking set $L = \{11, 22, 33, 44, 55\}$ between two adjacent slices.

It is convenient to express the λ 's in term of the following functions [60]

$$f_i(b, q) = \sum_{s=0}^{b-i} (-1)^s \binom{b-i}{s} (q - i - s)^{(b-i-s)}, \quad (2.3)$$

where $q^{(k)}$ is falling factorial defined as

$$q^{(k)} = \prod_{s=0}^{k-1} (q - s) . \quad (2.4)$$

Here we adopt the convention that $f_i(b, q) = 0$ if $b < i$.

For the case $b = 5$, one has

$$f_0(5, q) = q^5 - 15q^4 + 95q^3 - 325q^2 + 609q - 501 \quad (2.5)$$

$$f_1(5, q) = q^4 - 14q^3 + 77q^2 - 200q + 209 \quad (2.6)$$

$$f_2(5, q) = q^3 - 12q^2 + 50q - 73 \quad (2.7)$$

$$f_3(5, q) = q^2 - 9q + 21 \quad (2.8)$$

$$f_4(5, q) = q - 5 \quad (2.9)$$

$$f_5(5, q) = 1 , \quad (2.10)$$

and the chromatic polynomial for the $B_m(5)$ strip is

$$P(B_m(5), q) = \sum_{d=0}^5 \sum_{j=1}^{N_{B_m(5),d,\lambda}} c_{B_m(5),d,j}(q) c'_{B_m(5),d,j}(\lambda_{B_m(5),d,j}(q))^m , \quad (2.11)$$

where $N_{B_m(5),d,\lambda}$'s are equal to 1, 2, 5, 9, 9, 1 for d from 0 to 5. We find that the total number of distinct λ 's is

$$N_{P,B_m(5),\lambda} = \sum_{d=0}^5 N_{B_m(5),d,\lambda} = 27 . \quad (2.12)$$

We note that our result (2.12) differs from the pattern $N_{P,B_m(b),\lambda} = 2^b$ that was found for $1 \leq b \leq 4$, namely $N_{P,B_m(1),\lambda} = 2$, $N_{P,B_m(2),\lambda} = 2^2$ [9], $N_{P,B_m(3),\lambda} = 2^3$ [10], and $N_{P,B_m(4),\lambda} = 2^4$ [11].

We calculate that the eigenvalue with coefficient of degree 0 in q is unique and is given by

$$\lambda_{B_m(5),0,1} = f_0(5, q) = q^5 - 15q^4 + 95q^3 - 325q^2 + 609q - 501 . \quad (2.13)$$

For the eigenvalues with coefficients of degree 1 in q we find

$$\lambda_{B_m(5),1,1} = -f_1(5, q) + 4f_2(5, q) = -q^4 + 18q^3 - 125q^2 + 400q - 501 \quad (2.14)$$

$$\lambda_{B_m(5),1,2} = -f_1(5, q) - f_2(5, q) = -(q - 4)(q^3 - 9q^2 + 29q - 34) . \quad (2.15)$$

Next, for the eigenvalues with coefficients of degree 2 in q we obtain

$$\lambda_{B_m(5),2,1} = f_2(5, q) - 6f_3(5, q) + 6f_4(5, q) = q^3 - 18q^2 + 110q - 229 \quad (2.16)$$

$$\lambda_{B_m(5),2,2} = f_2(5, q) - f_3(5, q) - 4f_4(5, q) = q^3 - 13q^2 + 55q - 74 \quad (2.17)$$

$$\lambda_{B_m(5),2,3} = f_2(5, q) + 2f_3(5, q) + 2f_4(5, q) = q^3 - 10q^2 + 34q - 41 \quad (2.18)$$

$$\lambda_{B_m(5),2,4} = f_2(5, q) - 3f_3(5, q) = q^3 - 15q^2 + 77q - 136 \quad (2.19)$$

$$\lambda_{B_m(5),2,5} = f_2(5, q) + 2f_3(5, q) = q^3 - 10q^2 + 32q - 31 . \quad (2.20)$$

For $1 \leq j \leq 3$ these $\lambda_{B_m(5),2,j}$'s correspond to the partition [2] and for $j = 4, 5$ to the partition [11].

Proceeding to the eigenvalues with coefficients of degree 3, we find

$$\lambda_{B_m(5),3,1} = -f_3(5, q) + 6f_4(5, q) - 6f_5(5, q) = -q^2 + 15q - 57 \quad (2.21)$$

$$\lambda_{B_m(5),3,2} = -f_3(5, q) + f_4(5, q) + 4f_5(5, q) = -q^2 + 10q - 22 \quad (2.22)$$

$$\lambda_{B_m(5),3,3} = -f_3(5, q) - 2f_4(5, q) - 2f_5(5, q) = -q^2 + 7q - 13 \quad (2.23)$$

$$\lambda_{B_m(5),3,4} = -f_3(5, q) + 4f_4(5, q) - 2f_5(5, q) = -q^2 + 13q - 43 \quad (2.24)$$

$$\lambda_{B_m(5),3,5} = -f_3(5, q) + f_4(5, q) + f_5(5, q) = -(q - 5)^2 \quad (2.25)$$

$$\lambda_{B_m(5),3,6} = -f_3(5, q) - f_4(5, q) + 3f_5(5, q) = -q^2 + 8q - 13 \quad (2.26)$$

$$\lambda_{B_m(5),3,7} = -f_3(5, q) - 3f_4(5, q) - 3f_5(5, q) = -(q - 3)^2 \quad (2.27)$$

$$\lambda_{B_m(5),3,8} = -f_3(5, q) + 2f_4(5, q) = -q^2 + 11q - 31 \quad (2.28)$$

$$\lambda_{B_m(5),3,9} = -f_3(5, q) - 3f_4(5, q) = -q^2 + 6q - 6 . \quad (2.29)$$

Here the $\lambda_{B_m(5),3,j}$ for $1 \leq j \leq 3$ correspond to the partition [3], for $4 \leq j \leq 7$ to the partition [21], and for $j = 8, 9$ to the partition [111].

For the eigenvalues with coefficients of degree 4 in q , we calculate

$$\lambda_{B_m(5),4,1} = f_4(5, q) - 4f_5(5, q) = q - 9 \quad (2.30)$$

$$\lambda_{B_m(5),4,2} = f_4(5, q) + f_5(5, q) = q - 4 \quad (2.31)$$

$$\lambda_{B_m(5),4,3} = f_4(5, q) - 3f_5(5, q) = q - 8 \quad (2.32)$$

$$\lambda_{B_m(5),4,4} = f_4(5, q) + 3f_5(5, q) = q - 2 \quad (2.33)$$

$$\lambda_{B_m(5),4,5} = f_4(5, q) + 2f_5(5, q) = q - 3 \quad (2.34)$$

$$\lambda_{B_m(5),4,6} = f_4(5, q) = q - 5 \quad (2.35)$$

$$\lambda_{B_m(5),4,7} = f_4(5, q) - 2f_5(5, q) = q - 7 \quad (2.36)$$

$$\lambda_{B_m(5),4,8} = f_4(5, q) - f_5(5, q) = q - 6 \quad (2.37)$$

$$\lambda_{B_m(5),4,9} = f_4(5, q) + 4f_5(5, q) = q - 1 . \quad (2.38)$$

Here the $\lambda_{B_m(5),4,j}$ for $j = 1, 2$ correspond to the partition [4], for $j = 3, 5, 6$ to the partition [31], for $j = 4, 6, 7$ to the partition [211], for $j = 5, 7$ to the partition [22], and for $j = 8, 9$ to the partition [1111]. Notice that the terms $\lambda_{B_m(5),4,j}$ for $5 \leq j \leq 7$ involve degeneracies from two different associated partitions.

Finally, for the eigenvalue with coefficient of degree 5, we have

$$\lambda_{B_m(5),5,1} = -f_5(5, q) = -1 . \quad (2.39)$$

This involves a degeneracy from all the possible partitions.

The coefficient $c_{B_m(5),d,j}(q)c'_{B_m(5),d,j}$ of degree 0 in q is

$$c_{B_m(5),0,1}c'_{B_m(5),0,1} = 1 . \quad (2.40)$$

The coefficients of degree 1 in q are

$$c_{B_m(5),1,1}c'_{B_m(5),1,1} = q - 1 \quad (2.41)$$

$$c_{B_m(5),1,2}c'_{B_m(5),1,2} = 4(q - 1) , \quad (2.42)$$

where $c_{B_m(5),1,j} = q - 1$ for $j = 1, 2$, and $c'_{B_m(5),1,j} = 1, 4$ for $j = 1, 2$.

For $d = 2$, there are partitions $[2]$ and $[11]$, and the associated $c_{X,2}$ are $c_{[2],2} = \frac{1}{2}q(q - 3)$ and $c_{[11],2} = \frac{1}{2}(q - 1)(q - 2)$. The coefficients of degree 2 in q are listed in Table I

TABLE I. Coefficients of degree 2 in q for $B_m(5)$ strip.

j	partition	$c'_{B_m(5),2,j}$	$c_{B_m(5),2,j}c'_{B_m(5),2,j}$
1	$[2]$	1	$\frac{1}{2}q(q - 3)$
2	$[2]$	4	$2q(q - 3)$
3	$[2]$	5	$\frac{5}{2}q(q - 3)$
4	$[11]$	4	$2(q - 1)(q - 2)$
5	$[11]$	6	$3(q - 1)(q - 2)$

For $d = 3$, there are partitions $[3]$, $[21]$ and $[111]$, and the associated $c_{X,3}$ are $c_{[3],3} = \frac{1}{6}q(q - 1)(q - 5)$, $c_{[21],3} = \frac{1}{6}q(q - 2)(q - 4)$ and $c_{[111],3} = \frac{1}{6}(q - 1)(q - 2)(q - 3)$. The coefficients of degree 3 in q are listed in Table II

TABLE II. Coefficients of degree 3 in q for $B_m(5)$ strip.

j	partition	$c'_{B_m(5),3,j}$	$c_{B_m(5),3,j}c'_{B_m(5),3,j}$
1	$[3]$	1	$\frac{1}{6}q(q - 1)(q - 5)$
2	$[3]$	4	$\frac{2}{3}q(q - 1)(q - 5)$
3	$[3]$	5	$\frac{5}{6}q(q - 1)(q - 5)$
4	$[21]$	8	$\frac{4}{3}q(q - 2)(q - 4)$
5	$[21]$	10	$\frac{5}{3}q(q - 2)(q - 4)$
6	$[21]$	12	$2q(q - 2)(q - 4)$
7	$[21]$	10	$\frac{5}{3}q(q - 2)(q - 4)$
8	$[111]$	6	$(q - 1)(q - 2)(q - 3)$
9	$[111]$	4	$\frac{2}{3}(q - 1)(q - 2)(q - 3)$

For $d = 4$, there are partitions $[4]$, $[31]$, $[22]$, $[211]$ and $[1111]$, and the associated $c_{X,4}$ are $c_{[4],4} = \frac{1}{24}q(q-1)(q-2)(q-7)$, $c_{[31],4} = \frac{1}{24}q(q-1)(q-3)(q-6)$, $c_{[22],4} = \frac{1}{24}q(q-1)(q-4)(q-5)$, $c_{[211],4} = \frac{1}{24}q(q-2)(q-3)(q-5)$, and $c_{[1111],4} = \frac{1}{24}(q-1)(q-2)(q-3)(q-4)$. The coefficients of degree 4 in q are listed in Table III

TABLE III. Coefficients of degree 4 in q for $B_m(5)$ strip.

j	partition	$c'_{B_m(5),4,j}$	$c_{B_m(5),4,j}c'_{B_m(5),4,j}$
1	$[4]$	1	$\frac{1}{24}q(q-1)(q-2)(q-7)$
2	$[4]$	4	$\frac{1}{6}q(q-1)(q-2)(q-7)$
3	$[31]$	12	$\frac{1}{2}q(q-1)(q-3)(q-6)$
4	$[211]$	12	$\frac{1}{2}q(q-2)(q-3)(q-5)$
5	$[31]$	18	$\frac{1}{6}q(q-1)(7q^2 - 63q + 131)$
	$[22]$	10	
6	$[31]$	15	$\frac{5}{4}q(q-3)(q^2 - 7q + 8)$
	$[211]$	15	
7	$[211]$	18	$\frac{1}{6}q(q-5)(7q^2 - 35q + 37)$
	$[22]$	10	
8	$[1111]$	4	$\frac{1}{6}(q-1)(q-2)(q-3)(q-4)$
9	$[1111]$	1	$\frac{1}{24}(q-1)(q-2)(q-3)(q-4)$

Finally, the coefficient of degree 5 in q is given by

$$c_{B_m(5),5,1}c'_{B_m(5),5,1} = q^5 - 15q^4 + 75q^3 - 145q^2 + 89q - 1 . \quad (2.43)$$

The sum of all the coefficients is equal to

$$\begin{aligned} C(B_m(5), q) &= \sum_{d=0}^5 \sum_{j=1}^{N_{B_m(5),d,\lambda}} c_{B_m(5),d,j}(q) c'_{B_m(5),d,j} \\ &= q(q-1)(q-2)(q-3)(q-4) = P(K_5, q) . \end{aligned} \quad (2.44)$$

The chromatic number is $\chi(B_m(5)) = 5$.

The locus $\mathcal{B}$ and chromatic zeros for the $B_m(5)$ strip with $L_x = m = 20$ are shown in Fig. 1. The locus $\mathcal{B}$ crosses the real q -axis at $q = 0, 2, 4$ and $q_c(B_m(5))$, where

$$q_c(B_m(5)) \simeq 4.50634 \dots . \quad (2.45)$$




FIG. 1. Locus $\mathcal{B}$ for the $m \rightarrow \infty$ limit of the family $B_m(5)$ with toroidal boundary conditions and chromatic zeros for $B_m(5)$ with $m = 20$ (i.e., $n = 100$).

The locus $\mathcal{B}$ has support for $\text{Re}(q) \geq 0$, and separates the q plane into four regions. The outermost one, region R_1 , extends to infinite $|q|$ and includes the intervals $q \geq q_c(B_m(5))$ and $q \leq 0$ on the real q axis. Region R_2 includes the real interval $4 \leq q \leq q_c(B_m(5))$, region R_3 includes the real interval $2 \leq q \leq 4$, while region R_4 includes the real interval $0 \leq q \leq 2$. In regions R_i , $1 \leq i \leq 4$, the dominant terms are $\lambda_{B_m(5),0,1}$, $\lambda_{B_m(5),3,1}$, $\lambda_{B_m(5),2,1}$, and $\lambda_{B_m(5),1,1}$, respectively. Thus, the $q_c(B_m(5))$ given in (2.45) is the degeneracy between $|\lambda_{B_m(5),0,1}|$ and $|\lambda_{B_m(5),3,1}|$, and is the real solution of $q^5 - 15q^4 + 95q^3 - 326q^2 + 624q - 558 = 0$.

B. $L_y = 6$ toroidal strip

We have also succeeded in obtaining the chromatic polynomial for $B_m(6)$, i.e., the $6 \times L_x$ strips with each transverse slice forming a K_6 graph, and with edge linking set $L = \{11, 22, 33, 44, 55, 66\}$ between two adjacent slices.

For $b = 6$, one has

$$f_0(6, q) = q^6 - 21q^5 + 190q^4 - 965q^3 + 2944q^2 - 5155q + 4051 \quad (2.46)$$

$$f_1(6, q) = q^5 - 20q^4 + 165q^3 - 710q^2 + 1609q - 1546 \quad (2.47)$$

$$f_2(6, q) = q^4 - 18q^3 + 125q^2 - 400q + 501 \quad (2.48)$$

$$f_3(6, q) = q^3 - 15q^2 + 77q - 136 \quad (2.49)$$

$$f_4(6, q) = q^2 - 11q + 31 \quad (2.50)$$

$$f_5(6, q) = q - 6 \quad (2.51)$$

$$f_6(6, q) = 1, \quad (2.52)$$

and the chromatic polynomial for the $B_m(6)$ strip is

$$P((B_m(6), q) = \sum_{d=0}^6 \sum_{j=1}^{N_{B_m(6),d,\lambda}} c_{B_m(6),d,j}(q) c'_{B_m(6),d,j}(\lambda_{B_m(6),d,j}(q))^m, \quad (2.53)$$

where $N_{B_m(6),d,\lambda}$'s are equal to 1, 2, 5, 10, 16, 11, 1 for d from 0 to 6. We find that the total number of distinct λ 's is

$$N_{P,B_m(6),\lambda} = 46. \quad (2.54)$$

The eigenvalue with coefficient of degree 0 in q is unique and is given by

$$\lambda_{B_m(6),0,1} = f_0(6, q) = q^6 - 21q^5 + 190q^4 - 965q^3 + 2944q^2 - 5155q + 4051 . \quad (2.55)$$

For the eigenvalues with coefficients of degree 1 in q we have

$$\lambda_{B_m(6),1,1} = -f_1(6, q) + 5f_2(6, q) = -q^5 + 25q^4 - 255q^3 + 1335q^2 - 3609q + 4051 \quad (2.56)$$

$$\lambda_{B_m(6),1,2} = -f_1(6, q) - f_2(6, q) = -(q - 5)(q^4 - 14q^3 + 77q^2 - 200q + 209) . \quad (2.57)$$

For the eigenvalues with coefficients of degree 2 in q we find

$$\lambda_{B_m(6),2,1} = f_2(6, q) - 8f_3(6, q) + 12f_4(6, q) = q^4 - 26q^3 + 257q^2 - 1148q + 1961 \quad (2.58)$$

$$\lambda_{B_m(6),2,2} = f_2(6, q) - 2f_3(6, q) - 6f_4(6, q) = q^4 - 20q^3 + 149q^2 - 488q + 587 \quad (2.59)$$

$$\lambda_{B_m(6),2,3} = f_2(6, q) + 2f_3(6, q) + 2f_4(6, q) = q^4 - 16q^3 + 97q^2 - 268q + 291 \quad (2.60)$$

$$\lambda_{B_m(6),2,4} = f_2(6, q) - 4f_3(6, q) = q^4 - 22q^3 + 185q^2 - 708q + 1045 \quad (2.61)$$

$$\lambda_{B_m(6),2,5} = f_2(6, q) + 2f_3(6, q) = q^4 - 16q^3 + 95q^2 - 246q + 229 . \quad (2.62)$$

For $1 \leq j \leq 3$, these $\lambda_{B_m(6),2,j}$ correspond to the partition [2] and for $j = 4, 5$ to the partition [11].

For the eigenvalues with coefficients of degree 3 in q we obtain

$$\lambda_{B_m(6),3,1} = -f_3(6, q) + 9f_4(6, q) - 18f_5(6, q) + 6f_6(6, q) = -q^3 + 24q^2 - 194q + 529 \quad (2.63)$$

$$\lambda_{B_m(6),3,2} = -f_3(6, q) + 3f_4(6, q) + 6f_5(6, q) - 6f_6(6, q) = -q^3 + 18q^2 - 104q + 187 \quad (2.64)$$

$$\lambda_{B_m(6),3,3} = -f_3(6, q) - f_4(6, q) + 2f_5(6, q) + 6f_6(6, q) = -q^3 + 14q^2 - 64q + 99 \quad (2.65)$$

$$\lambda_{B_m(6),3,4} = -f_3(6, q) - 3f_4(6, q) - 6f_5(6, q) - 6f_6(6, q) = -q^3 + 12q^2 - 50q + 73 \quad (2.66)$$

$$\lambda_{B_m(6),3,5} = -f_3(6, q) + 6f_4(6, q) - 6f_5(6, q) = -q^3 + 21q^2 - 149q + 358 \quad (2.67)$$

$$\lambda_{B_m(6),3,6} = -f_3(6, q) + 2f_4(6, q) + 2f_5(6, q) = -(q - 6)(q^2 - 11q + 31) \quad (2.68)$$

$$\lambda_{B_m(6),3,7} = -f_3(6, q) + 6f_5(6, q) = -q^3 + 15q^2 - 71q + 100 \quad (2.69)$$

$$\lambda_{B_m(6),3,8} = -f_3(6, q) - 3f_4(6, q) - 3f_5(6, q) = -q^3 + 12q^2 - 47q + 61 \quad (2.70)$$

$$\lambda_{B_m(6),3,9} = -f_3(6, q) + 3f_4(6, q) = -q^3 + 18q^2 - 110q + 229 \quad (2.71)$$

$$\lambda_{B_m(6),3,10} = -f_3(6, q) - 3f_4(6, q) = -q^3 + 12q^2 - 44q + 43 . \quad (2.72)$$

Here the $\lambda_{B_m(6),3,j}$ for $1 \leq j \leq 4$ correspond to the partition $[3]$, for $5 \leq j \leq 8$ to the partition $[21]$, and for $j = 9, 10$ to the partition $[111]$.

Proceeding to the eigenvalues with coefficients of degree 4 in q we calculate

$$\lambda_{B_m(6),4,1} = f_4(6, q) - 8f_5(6, q) + 12f_6(6, q) = q^2 - 19q + 91 \quad (2.73)$$

$$\lambda_{B_m(6),4,2} = f_4(6, q) - 2f_5(6, q) - 6f_6(6, q) = q^2 - 13q + 37 \quad (2.74)$$

$$\lambda_{B_m(6),4,3} = f_4(6, q) + 2f_5(6, q) + 2f_6(6, q) = q^2 - 9q + 21 \quad (2.75)$$

$$\lambda_{B_m(6),4,4} = f_4(6, q) - 6f_6(6, q) = q^2 - 11q + 25 \quad (2.76)$$

$$\lambda_{B_m(6),4,5} = f_4(6, q) - 2f_5(6, q) - 2f_6(6, q) = q^2 - 13q + 41 \quad (2.77)$$

$$\lambda_{B_m(6),4,6} = f_4(6, q) = q^2 - 11q + 31 \quad (2.78)$$

$$\lambda_{B_m(6),4,7} = f_4(6, q) + 3f_5(6, q) + 3f_6(6, q) = (q - 4)^2 \quad (2.79)$$

$$\lambda_{B_m(6),4,8} = f_4(6, q) - 6f_5(6, q) + 6f_6(6, q) = q^2 - 17q + 73 \quad (2.80)$$

$$\lambda_{B_m(6),4,9} = f_4(6, q) + f_5(6, q) - 3f_6(6, q) = q^2 - 10q + 22 \quad (2.81)$$

$$\lambda_{B_m(6),4,10} = f_4(6, q) + 4f_5(6, q) + 6f_6(6, q) = q^2 - 7q + 13 \quad (2.82)$$

$$\lambda_{B_m(6),4,11} = f_4(6, q) - 4f_5(6, q) + 2f_6(6, q) = q^2 - 15q + 57 \quad (2.83)$$

$$\lambda_{B_m(6),4,12} = f_4(6, q) + 2f_5(6, q) - 4f_6(6, q) = q^2 - 9q + 15 \quad (2.84)$$

$$\lambda_{B_m(6),4,13} = f_4(6, q) - f_5(6, q) - f_6(6, q) = (q - 6)^2 \quad (2.85)$$

$$\lambda_{B_m(6),4,14} = f_4(6, q) + 4f_5(6, q) + 4f_6(6, q) = q^2 - 7q + 11 \quad (2.86)$$

$$\lambda_{B_m(6),4,15} = f_4(6, q) - 2f_5(6, q) = q^2 - 13q + 43 \quad (2.87)$$

$$\lambda_{B_m(6),4,16} = f_4(6, q) + 4f_5(6, q) = q^2 - 7q + 7 . \quad (2.88)$$

Here the $\lambda_{B_m(6),4,j}$ for $1 \leq j \leq 3$ correspond to the partition [4], for $4 \leq j \leq 8$ to the partition [31], for $9 \leq j \leq 11$ to the partition [22], for $11 \leq j \leq 14$ to the partition [211], and for $j = 15, 16$ to the partition [1111]. Notice that the term $\lambda_{B_m(6),4,11}$ involves a degeneracy from two different associated partitions.

For the eigenvalues with coefficients of degree 5 in q we find

$$\lambda_{B_m(6),5,1} = -f_5(6, q) + 5f_6(6, q) = -q + 11 \quad (2.89)$$

$$\lambda_{B_m(6),5,2} = -f_5(6, q) - f_6(6, q) = -q + 5 \quad (2.90)$$

$$\lambda_{B_m(6),5,3} = -f_5(6, q) + 4f_6(6, q) = -q + 10 \quad (2.91)$$

$$\lambda_{B_m(6),5,4} = -f_5(6, q) = -q + 6 \quad (2.92)$$

$$\lambda_{B_m(6),5,5} = -f_5(6, q) - 2f_6(6, q) = -q + 4 \quad (2.93)$$

$$\lambda_{B_m(6),5,6} = -f_5(6, q) + 3f_6(6, q) = -q + 9 \quad (2.94)$$

$$\lambda_{B_m(6),5,7} = -f_5(6, q) + f_6(6, q) = -q + 7 \quad (2.95)$$

$$\lambda_{B_m(6),5,8} = -f_5(6, q) - 3f_6(6, q) = -q + 3 \quad (2.96)$$

$$\lambda_{B_m(6),5,9} = -f_5(6, q) + 2f_6(6, q) = -q + 8 \quad (2.97)$$

$$\lambda_{B_m(6),5,10} = -f_5(6, q) - 4f_6(6, q) = -q + 2 \quad (2.98)$$

$$\lambda_{B_m(6),5,11} = -f_5(6, q) - 5f_6(6, q) = -q + 1 . \quad (2.99)$$

Here $\lambda_{B_m(6),5,j}$ for $j = 1, 2$ correspond to the partition [5], for $3 \leq j \leq 5$ to the partition [41], for $5 \leq j \leq 7$ to the partition [32], for $j = 4, 6, 8$ to the partition [311], for $j = 2, 8, 9$ to the partition [221], for $j = 4, 9, 10$ to the partition [2111], and for $j = 7, 11$ to the partition [11111]. Notice that the terms $\lambda_{B_m(6),5,j}$ for $j = 2$ and $5 \leq j \leq 9$ involve degeneracies from two different associated partitions, and for $j = 4$ from three associated partitions.

Finally, for the eigenvalue with coefficient of degree 6 in q we have

$$\lambda_{B_m(6),6,1} = f_6(6, q) = 1 . \quad (2.100)$$

This involves a degeneracy from all the possible partitions.

The coefficient $c_{B_m(6),d,j}(q)c'_{B_m(6),d,j}$ of degree 0 in q is

$$c_{B_m(6),0,1}c'_{B_m(6),0,1} = 1 . \quad (2.101)$$

The coefficients of degree 1 in q are

$$c_{B_m(6),1,1}c'_{B_m(6),1,1} = q - 1 \quad (2.102)$$

$$c_{B_m(6),1,2}c'_{B_m(6),1,2} = 5(q - 1) , \quad (2.103)$$

where $c_{B_m(6),1,j} = q - 1$ for $j = 1, 2$, and $c'_{B_m(6),1,j} = 1, 5$ for $j = 1, 2$.

The coefficients of degree 2 in q are listed in Table IV

TABLE IV. Coefficients of degree 2 in q for $B_m(6)$ strip.

j	partition	$c'_{B_m(6),2,j}$	$c_{B_m(6),2,j}c'_{B_m(6),2,j}$
1	[2]	1	$\frac{1}{2}q(q - 3)$
2	[2]	5	$\frac{5}{2}q(q - 3)$
3	[2]	9	$\frac{9}{2}q(q - 3)$
4	[11]	5	$\frac{5}{2}(q - 1)(q - 2)$
5	[11]	10	$5(q - 1)(q - 2)$

The coefficients of degree 3 in q are listed in Table V

TABLE V. Coefficients of degree 3 in q for $B_m(6)$ strip.

j	partition	$c'_{B_m(6),3,j}$	$c_{B_m(6),3,j} c'_{B_m(6),3,j}$
1	[3]	1	$\frac{1}{6}q(q-1)(q-5)$
2	[3]	5	$\frac{5}{6}q(q-1)(q-5)$
3	[3]	9	$\frac{3}{2}q(q-1)(q-5)$
4	[3]	5	$\frac{5}{6}q(q-1)(q-5)$
5	[21]	10	$\frac{5}{3}q(q-2)(q-4)$
6	[21]	18	$3q(q-2)(q-4)$
7	[21]	20	$\frac{10}{3}q(q-2)(q-4)$
8	[21]	32	$\frac{16}{3}q(q-2)(q-4)$
9	[111]	10	$\frac{5}{3}(q-1)(q-2)(q-3)$
10	[111]	10	$\frac{5}{3}(q-1)(q-2)(q-3)$

The coefficients of degree 4 in q are listed in Table VI

TABLE VI. Coefficients of degree 4 in q for $B_m(6)$ strip.

j	partition	$c'_{B_m(6),4,j}$	$c_{B_m(6),4,j} c'_{B_m(6),4,j}$
1	[4]	1	$\frac{1}{24}q(q-1)(q-2)(q-7)$
2	[4]	5	$\frac{5}{24}q(q-1)(q-2)(q-7)$
3	[4]	9	$\frac{3}{8}q(q-1)(q-2)(q-7)$
4	[31]	30	$\frac{5}{4}q(q-1)(q-3)(q-6)$
5	[31]	27	$\frac{9}{8}q(q-1)(q-3)(q-6)$
6	[31]	15	$\frac{5}{8}q(q-1)(q-3)(q-6)$
7	[31]	48	$2q(q-1)(q-3)(q-6)$
8	[31]	15	$\frac{5}{8}q(q-1)(q-3)(q-6)$
9	[22]	32	$\frac{4}{3}q(q-1)(q-4)(q-5)$
10	[22]	10	$\frac{5}{12}q(q-1)(q-4)(q-5)$
11	[22]	18	$\frac{1}{2}q(q-5)(2q-3)(2q-7)$
	[211]	30	
12	[211]	30	$\frac{5}{4}q(q-2)(q-3)(q-5)$
13	[211]	48	$2q(q-2)(q-3)(q-5)$
14	[211]	27	$\frac{9}{8}q(q-2)(q-3)(q-5)$
15	[1111]	10	$\frac{5}{12}(q-1)(q-2)(q-3)(q-4)$
16	[1111]	5	$\frac{5}{24}(q-1)(q-2)(q-3)(q-4)$

For $d = 5$, there are partitions $[5]$, $[41]$, $[32]$, $[311]$, $[221]$, $[2111]$ and $[11111]$, and the associated $c_{X,5}$ are $c_{[5],5} = \frac{1}{120}q(q-1)(q-2)(q-3)(q-9)$, $c_{[41],5} = \frac{1}{120}q(q-1)(q-2)(q-4)(q-8)$, $c_{[32],5} = \frac{1}{120}q(q-1)(q-2)(q-5)(q-7)$, $c_{[311],5} = \frac{1}{120}q(q-1)(q-3)(q-4)(q-7)$, $c_{[221],5} = \frac{1}{120}q(q-1)(q-3)(q-5)(q-6)$, $c_{[2111],5} = \frac{1}{120}q(q-2)(q-3)(q-4)(q-6)$, and $c_{[11111],5} = \frac{1}{120}(q-1)(q-2)(q-3)(q-4)(q-5)$. We list the coefficients of degree 5 in q in Table VII

TABLE VII. Coefficients of degree 5 in q for $B_m(6)$ strip.

j	partition	$c'_{B_m(6),5,j}$	$c_{B_m(6),5,j}c'_{B_m(6),5,j}$
1	[5]	1	$\frac{1}{120}q(q-1)(q-2)(q-3)(q-9)$
2	[5]	5	$\frac{1}{4}q(q-1)(q-3)(q-4)(q-7)$
	[221]	25	
3	[41]	20	$\frac{1}{6}q(q-1)(q-2)(q-4)(q-8)$
4	[41]	36	$\frac{1}{5}q(q-4)(7q^3 - 77q^2 + 217q - 162)$
	[311]	96	
	[2111]	36	
5	[41]	40	$q(q-1)(q-2)(q^2 - 12q + 34)$
	[32]	80	
6	[32]	45	$\frac{1}{8}q(q-1)(q-7)(7q^2 - 49q + 78)$
	[311]	60	
7	[32]	25	$\frac{1}{4}(q-1)(q-2)(q-5)(q^2 - 7q + 2)$
	[11111]	5	
8	[311]	60	$\frac{1}{8}q(q-1)(q-3)(7q^2 - 77q + 202)$
	[221]	45	
9	[221]	80	$q(q-3)(q-6)(q^2 - 6q + 6)$
	[2111]	40	
10	[2111]	20	$\frac{1}{6}q(q-2)(q-3)(q-4)(q-6)$
11	[11111]	1	$\frac{1}{120}(q-1)(q-2)(q-3)(q-4)(q-5)$

Finally, the coefficient of degree 6 is

$$c_{B_m(6),6,1}c'_{B_m(6),6,1} = q^6 - 21q^5 + 160q^4 - 545q^3 + 814q^2 - 415q + 1 . \quad (2.104)$$

The sum of all the coefficients is equal to

$$\begin{aligned} C(B_m(6), q) &= \sum_{d=0}^6 \sum_{j=1}^{N_{B_m(6),d,\lambda}} c_{B_m(6),d,j}(q) c'_{B_m(6),d,j} \\ &= q(q-1)(q-2)(q-3)(q-4)(q-5) = P(K_6, q) . \end{aligned} \quad (2.105)$$

The chromatic number is $\chi(B_m(6)) = 6$.

The locus $\mathcal{B}$ and chromatic zeros for the $B_m(6)$ strip with $L_x = m = 15$ are shown in Fig. 2. The locus $\mathcal{B}$ crosses the real q -axis at $q = 0, 2, 4$ and $q_c(B_m(6))$, where

$$q_c(B_m(6)) \simeq 5.3236 \dots . \quad (2.106)$$




FIG. 2. Locus $\mathcal{B}$ for the $m \rightarrow \infty$ limit of the family $B_m(6)$ with toroidal boundary conditions and chromatic zeros for $B_m(6)$ with $m = 15$ (i.e., $n = 90$).

The locus $\mathcal{B}$ has support for $\text{Re}(q) \geq 0$, and separates the q plane into four regions. The outermost one, region R_1 , extends to infinite $|q|$ and includes the intervals $q \geq q_c(B_m(6))$ and $q \leq 0$ on the real q axis. Region R_2 includes the real interval $4 \leq q \leq q_c(B_m(6))$, region R_3 includes the real interval $2 \leq q \leq 4$, while region R_4 includes the real interval $0 \leq q \leq 2$. In regions R_i , $1 \leq i \leq 4$, the dominant terms are $\lambda_{B_m(6),0,1}$, $\lambda_{B_m(6),3,1}$, $\lambda_{B_m(6),2,1}$, and $\lambda_{B_m(6),1,1}$, respectively. Thus, the $q_c(B_m(6))$ given in (2.106) is the degeneracy between $|\lambda_{B_m(6),0,1}|$ and $|\lambda_{B_m(6),3,1}|$, and is the real solution of $q^5 - 19q^4 + 152q^3 - 660q^2 + 1600q - 1761 = 0$.

III. FAMILY OF KLEIN BOTTLE STRIPS

Consider the graph with L_x set of K_b with $L_x - 1$ edge linking sets $\{11, 22, \dots, bb\}$ and one edge linking set $\{1b, 2(b-1), \dots, b1\}$. This is the Klein bottle strip and will be denoted as $\bar{B}_m(b)$. The chromatic polynomials has the same form as given in eq. (2.1), and the λ 's are the same as those of the corresponding toroidal strip with the same b . The only difference is the coefficients.

A. $L_y = 5$ Klein bottle strip

For $b = 5$ considered here, the coefficients $c_{\bar{B}_m(5),d,j}(q)c'_{\bar{B}_m(5),d,j}$ are summarized in Table VIII

TABLE VIII. Coefficients for $\bar{B}_m(5)$ strip.

d	j	partition	$c'_{\bar{B}_m(5),d,j}$	$c_{\bar{B}_m(5),d,j}c'_{\bar{B}_m(5),d,j}$
0	1	[0]	1	1
1	1	[1]	1	$q - 1$
1	2	[1]	0	0
2	1	[2]	1	$\frac{1}{2}q(q - 3)$
2	2	[2]	0	0
2	3	[2]	1	$\frac{1}{2}q(q - 3)$
2	4	[11]	0	0
2	5	[11]	-2	$-(q - 1)(q - 2)$

3	1	[3]	1	$\frac{1}{6}q(q-1)(q-5)$
3	2	[3]	0	0
3	3	[3]	1	$\frac{1}{6}q(q-1)(q-5)$
3	4	[21]	0	0
3	5	[21]	2	$\frac{1}{3}q(q-2)(q-4)$
3	6	[21]	-4	$-\frac{2}{3}q(q-2)(q-4)$
3	7	[21]	2	$\frac{1}{3}q(q-2)(q-4)$
3	8	[111]	-2	$-\frac{1}{3}(q-1)(q-2)(q-3)$
3	9	[111]	0	0
4	1	[4]	1	$\frac{1}{24}q(q-1)(q-2)(q-7)$
4	2	[4]	0	0
4	3	[31]	0	0
4	4	[211]	0	0
4	5	[31]	-6	$-\frac{1}{6}q(q-1)(q^2-9q+17)$
		[22]	2	
4	6	[31]	3	$\frac{1}{4}q(q-3)(q^2-7q+8)$
		[211]	3	
4	7	[211]	-6	$-\frac{1}{6}q(q-5)(q^2-5q+7)$
		[22]	2	
4	8	[1111]	0	0
4	9	[1111]	1	$\frac{1}{24}(q-1)(q-2)(q-3)(q-4)$
5	1	see text		$q-1$

where for $d = 5$, it involves all possible partitions with appropriate $c'_{\bar{B}_m(5),5,j}$ which are not listed here. Notice that the coefficients for $\lambda_{\bar{B}_m(5),d,j}$ with $(d,j) = (1,2), (2,2), (2,4), (3,2), (3,4), (3,9), (4,2), (4,3), (4,4), (4,8)$ are zero, i.e., these ten λ 's do not contribute to the chromatic polynomial of the Klein bottle strip. We thus find that the total number of distinct λ 's for the $b = 5$ Klein bottle strip with indicated linking is

$$N_{P,\bar{B}_m(5),\lambda} = 17 . \quad (3.1)$$

The sum of all of the coefficients for the Klein bottle strip is

$$C(\bar{B}_m(5), q) = \sum_{d=0}^5 \sum_{j=1}^{N_{\bar{B}_m(5),d,\lambda}} c_{\bar{B}_m(5),d,j}(q) c'_{\bar{B}_m(5),d,j} = 0 , \quad (3.2)$$

which is easy to understand by the coloring argument.

B. $L_y = 6$ Klein bottle strip

For $b = 6$, the coefficients $c_{\bar{B}_m(6),d,j}(q)c'_{\bar{B}_m(6),d,j}$ are summarized in Table IX

TABLE IX. Coefficients for $\bar{B}_m(6)$ strip.

d	j	partition	$c'_{\bar{B}_m(6),d,j}$	$c_{\bar{B}_m(6),d,j}c'_{\bar{B}_m(6),d,j}$
0	1	[0]	1	1
1	1	[1]	1	$q - 1$
1	2	[1]	-1	$-(q - 1)$
2	1	[2]	1	$\frac{1}{2}q(q - 3)$
2	2	[2]	-1	$-\frac{1}{2}q(q - 3)$
2	3	[2]	3	$\frac{3}{2}q(q - 3)$
2	4	[11]	-1	$-\frac{1}{2}(q - 1)(q - 2)$
2	5	[11]	-2	$-(q - 1)(q - 2)$
3	1	[3]	1	$\frac{1}{6}q(q - 1)(q - 5)$
3	2	[3]	-1	$-\frac{1}{6}q(q - 1)(q - 5)$
3	3	[3]	3	$\frac{1}{2}q(q - 1)(q - 5)$
3	4	[3]	-3	$-\frac{1}{2}q(q - 1)(q - 5)$
3	5	[21]	-2	$-\frac{1}{3}q(q - 2)(q - 4)$
3	6	[21]	6	$q(q - 2)(q - 4)$
3	7	[21]	-4	$-\frac{2}{3}q(q - 2)(q - 4)$
3	8	[21]	0	0
3	9	[111]	-2	$-\frac{1}{3}(q - 1)(q - 2)(q - 3)$
3	10	[111]	2	$\frac{1}{3}(q - 1)(q - 2)(q - 3)$
4	1	[4]	1	$\frac{1}{24}q(q - 1)(q - 2)(q - 7)$
4	2	[4]	-1	$-\frac{1}{24}q(q - 1)(q - 2)(q - 7)$
4	3	[4]	3	$\frac{1}{8}q(q - 1)(q - 2)(q - 7)$
4	4	[31]	-6	$-\frac{1}{4}q(q - 1)(q - 3)(q - 6)$
4	5	[31]	9	$\frac{3}{8}q(q - 1)(q - 3)(q - 6)$
4	6	[31]	-9	$-\frac{3}{8}q(q - 1)(q - 3)(q - 6)$
4	7	[31]	0	0
4	8	[31]	-3	$-\frac{1}{8}q(q - 1)(q - 3)(q - 6)$
4	9	[22]	0	0
4	10	[22]	6	$\frac{1}{4}q(q - 1)(q - 4)(q - 5)$
4	11	[22]	6	$-\frac{1}{2}q(q - 5)$

		[211]	-6	
4	12	[211]	6	$\frac{1}{4}q(q-2)(q-3)(q-5)$
4	13	[211]	0	0
4	14	[211]	-9	$-\frac{3}{8}q(q-2)(q-3)(q-5)$
4	15	[1111]	2	$\frac{1}{12}(q-1)(q-2)(q-3)(q-4)$
4	16	[1111]	1	$\frac{1}{24}(q-1)(q-2)(q-3)(q-4)$
5	1	[5]	1	$\frac{1}{120}q(q-1)(q-2)(q-3)(q-9)$
5	2	[5]	-1	$\frac{1}{60}q(q-1)(q-3)(7q^2-77q+216)$
		[221]	15	
5	3	[41]	-4	$-\frac{1}{30}q(q-1)(q-2)(q-4)(q-8)$
5	4	[41]	12	$-q(q-2)(q-4)$
		[2111]	-12	
5	5	[41]	-8	$-\frac{1}{15}q(q-1)(q-2)(q-4)(q-8)$
5	6	[32]	15	$\frac{1}{40}q(q-1)(q-7)(q^2-7q+2)$
		[311]	-12	
5	7	[32]	-15	$-\frac{1}{60}(q-1)(q-2)(q-5)(7q^2-49q-6)$
		[11111]	1	
5	8	[311]	12	$-\frac{1}{40}q(q-1)(q-3)(q^2-11q+38)$
		[221]	-15	
5	9	[2111]	8	$\frac{1}{15}q(q-2)(q-3)(q-4)(q-6)$
5	10	[2111]	4	$\frac{1}{30}q(q-2)(q-3)(q-4)(q-6)$
5	11	[11111]	-1	$-\frac{1}{120}(q-1)(q-2)(q-3)(q-4)(q-5)$
6	1	see text		-1

where for $d = 6$, it involves all possible partitions with appropriate $c'_{\bar{B}_m(6),6,j}$ which are not listed here. Notice that the coefficients for $\lambda_{\bar{B}_m(6),d,j}$ with $(d, j) = (3, 8), (4, 7), (4, 9), (4, 13)$ are zero, i.e., these four λ 's do not contribute to the chromatic polynomial of the Klein bottle strip. We thus find that the total number of distinct λ 's for the $b = 6$ Klein bottle strip with indicated linking is

$$N_{P,\bar{B}_m(6)} = 42 . \quad (3.3)$$

The sum of all of the coefficients for the Klein bottle strip is again zero,

$$C(\bar{B}_m(6), q) = \sum_{d=0}^6 \sum_{j=1}^{N_{\bar{B}_m(6),d,\lambda}} c_{\bar{B}_m(6),d,j}(q) c'_{\bar{B}_m(6),d,j} = 0 . \quad (3.4)$$

IV. PROPERTIES OF EIGENVALUES, COEFFICIENTS, AND $\mathcal{B}$

We first list the number of eigenvalues, $N_{B_m(b),d,\lambda}$, for $1 \leq b \leq 6$, $0 \leq d \leq b$, the total number of eigenvalues, $N_{P,B_m(b),\lambda}$, and the points at which $\mathcal{B}$ crosses the real q axis, as well as the corresponding values for the Klein bottle strips in Table X.

TABLE X. Properties of P and $\mathcal{B}$ for strip graphs $B_m(b)$ and $\bar{B}_m(b)$. The properties apply for a given strip of size $L_y \times L_x$; some apply for arbitrary L_x , such as $N_{P,B_m(b),\lambda}$, while others apply for the infinite-length limit, such as the properties of the locus $\mathcal{B}$. For the boundary conditions in the y and x directions (BC_y , BC_x), P, and T denote periodic, and orientation-reversed (twisted) periodic. $N_{B_m(b),d,\lambda}$'s are abbreviated as $N_{d,\lambda}$, $N_{P,B_m(b),\lambda}$ is abbreviated as $N_{P,\lambda}$, and similarly for $\bar{B}_m(b)$ strips. The blank entries are zero. The column denoted BCR lists the points at which $\mathcal{B}$ crosses the real q axis; the largest of these is $q_c(B_m(b))$ for the given family $B_m(b)$ or $\bar{B}_m(b)$. Column labelled “SN” refers to whether $\mathcal{B}$ has support for negative $Re(q)$, indicated as yes (y) or no (n).

b	BC_y	BC_x	$N_{0,\lambda}$	$N_{1,\lambda}$	$N_{2,\lambda}$	$N_{3,\lambda}$	$N_{4,\lambda}$	$N_{5,\lambda}$	$N_{6,\lambda}$	$N_{P,\lambda}$	BCR	SN
1	P	P	1	1						2	0, 2	n
2	P	P	1	2	1					4	0, 2	n
2	P	TP	1	2	1					4	0, 2	n
3	P	P	1	2	4	1				8	0, 2, 3	n
3	P	TP	1	1	2	1				5	0, 2, 3	n
4	P	P	1	2	5	7	1			16	0, 2, 3.67	n
4	P	TP	1	2	5	7	1			16	0, 2, 3.67	n
5	P	P	1	2	5	9	9	1		27	0, 2, 4, 4.51	n
5	P	TP	1	1	3	6	5	1		17	0, 2, 4, 4.51	n
6	P	P	1	2	5	10	16	11	1	46	0, 2, 4, 5.32	n
6	P	TP	1	2	5	9	13	11	1	42	0, 2, 4, 5.32	n

We conjecture the following q value at which $\mathcal{B}$ crosses the real axis for the $B_m(b)$ strips,

$$\mathcal{B} \supset \{q = 0, 2, 4, \dots, 2[\frac{b-1}{2}], q_c(B_m(b))\} . \quad (4.1)$$

where $[\nu]$ denotes the integral part of ν , and the dominant eigenvalue in interval between $2(d' - 1)$ and $2d'$ is $\lambda_{B_m(b), d', 1}$ for $1 \leq d' \leq [\frac{b+1}{2}]$.

The upper and lower bounds for the eigenvalues with coefficients of degree $b - 1$ can be determined in the following theorem.

Theorem 1 The eigenvalues with coefficients of degree $b - 1$ are bounded between $(-1)^{b-1}(q - 1)$ and $(-1)^{b-1}(q - 2b + 1)$.

Proof. Denote the $b! \times b!$ matrix for the eigenvalues with coefficients of degree $b - 1$ as M_{b-1} . By theorem 2 of [58], the diagonal elements of M_{b-1} are all the same to be $(-1)^{b-1}f_{b-1}(b, q) = (-1)^{b-1}(q - b)$, and other non-zero elements of M_{b-1} are all the same to be $(-1)^b f_b(b, q) = (-1)^b$. Now consider $M'_{b-1} = M_{b-1} - (-1)^{b-1}(q - b)I$, where I is the $b! \times b!$ identity matrix. The number of non-zero elements, $(-1)^b$, in every row and columns of M'_{b-1} is $b - 1$ by the construction of the matrix. If we denote the eigenvalues of M'_{b-1} as λ' and the normalized eigenvector of M'_{b-1} as $W^T = (w_1, w_2, \dots, w_{b!})$, then we have

$$\begin{aligned} M'_{b-1}W &= \lambda'W \\ \Rightarrow |M'_{b-1}W|^2 &= |\lambda'W|^2 \\ \Rightarrow (b-1)(w_1^2 + w_2^2 + \dots w_{b!}^2) + 2 \sum_{i,j,i \neq j} w_i w_j &= \lambda'^2(w_1^2 + w_2^2 + \dots w_{b!}^2) \\ \Rightarrow \lambda'^2 &= b-1 + 2 \sum_{i,j,i \neq j} w_i w_j , \end{aligned} \quad (4.2)$$

where we sum all the possible multiplications of two different w_i and w_j in each row of M'_{b-1} . Therefore,

$$\begin{aligned} \lambda'^2 &\leq b-1 + \sum_{i,j,i \neq j} (w_i^2 + w_j^2) \\ &= b-1 + (b-1)(b-2) = (b-1)^2 , \end{aligned} \quad (4.3)$$

that is, $-(b-1) \leq \lambda' \leq b-1$. The upper and lower bounds for the eigenvalues of M_{b-1} are

$$(-1)^{b-1}(q - b) \pm (b-1) = \begin{cases} (-1)^{b-1}(q - 1) \\ (-1)^{b-1}(q - 2b + 1) \end{cases} . \quad (4.4)$$

It is easy to see that $(-1)^{b-1}(q - 2b + 1)$ is one of the principal terms and $(-1)^{b-1}(q - 1)$ is one of the alternating terms given in [58]. It follows that these values are realized as the largest and smallest values for the eigenvalues of M_{b-1} $\square$.

We shall show that the locus $\mathcal{B}$ crosses the real q axis at $q = 0$, and doesn't cross any $q \in \mathbb{R}_-$. We need the following lemma.

Lemma 1 On the non-positive real q axis, the dominant eigenvalue with coefficient of degree d is

$$\lambda_{B_m(b),d} = (-1)^d \sum_{j=0}^d (-1)^j \binom{d}{j} (b-d)^{(j)} f_{d+j}(b, q) . \quad (4.5)$$

Proof. The matrix for the eigenvalues with coefficients of degree d can be written as [58] $M_d = (-1)^d f_d(b, q)I + (-1)^{d+1} f_{d+1}(b, q)M_{d,1} + (-1)^{d+2} f_{d+2}(b, q)M_{d,2} + \dots$, where the non-zero elements are 1 in every $M_{d,j}$, and the number of these in each row and column is $\binom{d}{j}(b-d)^{(j)}$. By the same reason given in Theorem 1, the magnitudes of the eigenvalues of $M_{d,j}$ are bounded above by $\binom{d}{j}(b-d)^{(j)}$. Since the signs of $f_{d+j}(b, q)$ alternate for even j and odd j for any non-positive q value, $\lambda_{B_m(b),d}$ given in eq. (4.5) is the upper bound for the eigenvalues with coefficients of degree d . Since $\lambda_{B_m(b),d}$ is the eigenvalue with $c'_{B_m(b),d,j} = 1$ given in [58], the proof is completed. $\square$.

Theorem 2 The minimum real q at which $\mathcal{B}$ crosses the real q axis is $q = 0$.

Proof. We will begin with the proof that $|\lambda_{B_m(b),0}| > |\lambda_{B_m(b),1}| > \dots$ on the whole negative real q axis. The first few examples of $\lambda_{B_m(b),d}$'s are

$$\begin{aligned} \lambda_{B_m(b),0} &= f_0(b, q) \\ \lambda_{B_m(b),1} &= -f_1(b, q) + (b-1)f_2(b, q) \\ \lambda_{B_m(b),2} &= f_2(b, q) - 2(b-2)f_3(b, q) + (b-2)(b-3)f_4(b, q) \\ \lambda_{B_m(b),3} &= -f_3(b, q) + 3(b-3)f_4(b, q) - 3(b-3)(b-4)f_5(b, q) + (b-3)(b-4)(b-5)f_6(b, q) \\ \lambda_{B_m(b),4} &= f_4(b, q) - 4(b-4)f_5(b, q) + 6(b-4)(b-5)f_6(b, q) - 4(b-4)(b-5)(b-6)f_7(b, q) \\ &\quad + (b-4)(b-5)(b-6)(b-7)f_8(b, q) . \end{aligned} \quad (4.6)$$

To show that $f_0(b, q)$ has the largest magnitude on the negative real q axis, we only need $|f_i(b, q)| > |f_{i+1}(b, q) - (b-i-1)f_{i+2}(b, q)|$ as follows,

$$\begin{aligned}
& |f_i(b, q)| - |f_{i+1}(b, q) - (b - i - 1)f_{i+2}(b, q)| \\
&= (-1)^{b-i} f_i(b, q) - (-1)^{b-i-1} [f_{i+1}(b, q) - (b - i - 1)f_{i+2}(b, q)] \\
&= (-1)^{b-i} \left[\sum_{s=0}^{b-i} (-1)^s \binom{b-i}{s} (q - i - s)^{(b-i-s)} + f_{i+1}(b, q) - (b - i - 1)f_{i+2}(b, q) \right] \\
&= (-1)^{b-i} \left[(q - i)^{(b-i)} + \sum_{s=0}^{b-i-1} (-1)^{s+1} \binom{b-i}{s+1} (q - i - s - 1)^{(b-i-s-1)} \right. \\
&\quad \left. + \sum_{s=0}^{b-i-1} (-1)^s \binom{b-i-1}{s} (q - i - 1 - s)^{(b-i-1-s)} - (b - i - 1)f_{i+2}(b, q) \right] \\
&= (-1)^{b-i} \left[(q - i)^{(b-i)} + (-1)^{b-i} + (-1)^{b-i-1} + \sum_{s=0}^{b-i-2} (-1)^{s+1} (q - i - 1 - s)^{(b-i-1-s)} \right. \\
&\quad \left. \times \left(\frac{(b-i)!}{(s+1)!(b-i-s-1)!} - \frac{(b-i-1)!}{s!(b-i-1-s)!} \right) - (b - i - 1)f_{i+2}(b, q) \right] \\
&= (-1)^{b-i} \left[(q - i)^{(b-i)} + \sum_{s=0}^{b-i-2} (-1)^{s+1} (q - i - 1 - s)^{(b-i-1-s)} \binom{b-i-1}{s+1} \right. \\
&\quad \left. - (b - i - 1) \sum_{s=0}^{b-i-2} (-1)^s \binom{b-i-2}{s} (q - i - 2 - s)^{(b-i-2-s)} \right] \\
&= (-1)^{b-i} \left[(q - i)^{(b-i)} + \sum_{s=0}^{b-i-2} (-1)^{s+1} (q - i - 2 - s)^{(b-i-2-s)} \right. \\
&\quad \left. \times \left[(q - i - 1 - s) \frac{(b-i-1)!}{(s+1)!(b-i-2-s)!} + (b-i-1) \frac{(b-i-2)!}{s!(b-i-2-s)!} \right] \right] \\
&= (-1)^{b-i} \left[(q - i)^{(b-i)} + \sum_{s=0}^{b-i-2} (-1)^{s+1} (q - i - 2 - s)^{(b-i-2-s)} (b-i-1) \binom{b-i-2}{s} \frac{q-i}{s+1} \right] \\
&> 0 \quad \text{for } q < 0. \tag{4.7}
\end{aligned}$$

For $q = 0$, all above argument are still valid, with the exception that $|f_i(b, 0)| = |f_{i+1}(b, 0) - (b - i - 1)f_{i+2}(b, 0)|$ when $i = 0$, i.e. $|\lambda_{B_m(b), 0}| = |\lambda_{B_m(b), 1}|$ for $q = 0$. $\square$.

We will prove that the maximal value of q where $\mathcal{B}$ intersects the (positive) real axis is smaller than $q = b$ for $b \geq 2$. We have the following lemma.

Lemma 2 $f_i(b, q) \geq 0$ if $q \geq b$.

Proof. Since we have $(q - i - s)^{(b-i-s)} > 0$ when $q \geq b$, one only need to show that the magnitudes of the terms in eq. (2.3) decrease for s from 0 to $b - i$.

$$\begin{aligned}
& \binom{b-i}{s} (q-i-s)^{(b-i-s)} - \binom{b-i}{s+1} (q-i-s-1)^{(b-i-s-1)} \\
&= (q-i-s-1)^{(b-i-s-1)} \left[\frac{(b-i)!}{s!(b-i-s)!} (q-i-s) - \frac{(b-i)!}{(s+1)!(b-i-s-1)!} \right] \\
&= (q-i-s-1)^{(b-i-s-1)} \frac{(b-i)!}{(s+1)!(b-i-s)!} \left[(s+1)(q-i-s) - (b-i-s) \right] \\
&= (q-i-s-1)^{(b-i-s-1)} \frac{(b-i)!}{(s+1)!(b-i-s)!} \left[s(q-i-s) + q-b \right] \\
&\geq 0 \quad \text{for } 0 \leq s \leq b-i-1 \text{ and } q \geq b, \tag{4.8}
\end{aligned}$$

where the equal sign is realized only when $s = 0$ and $q = b$. Therefore, $f_i(b, q) = 0$ only when $i = b - 1$ at $q = b$, and $f_i(b, q) > 0$ for any other $0 \leq i \leq b$ and $q \geq b$. $\square$.

It is known that $q_c(B_m(1)) = 2$, $q_c(B_m(2)) = 2$ [9,6], and $q_c(B_m(3)) = 3$ [10]. Thus, for $b = 1$, $q_c = b + 1$ while for $b = 2$ and for $b = 3$, q_c has the respective values $q_c = b$. It is of interest to ask how $q_c(B_m(b))$ is related to b for $b \geq 4$. We determine this relation with the following theorem.

Theorem 3 $q_c(B_m(b)) < b$ for $b \geq 4$.

Proof. By a reason similar to that given in Lemma 1 and the result of Lemma 2, the magnitude of the eigenvalues with coefficients of degree d are bounded above by

$$\lambda'_{B_m(b),d} = \sum_{j=0}^d \binom{d}{j} (b-d)^{(j)} f_{d+j}(b, q) \quad \text{for } q \geq b, \tag{4.9}$$

where it is equal to $\lambda_{B_m(b),b-1,1}$ for $d = b - 1$ at $q = b$ and equal to $\lambda_{B_m(b),0,1}$ for $d = 0$ and any q . The first few examples of $\lambda'_{B_m(b),d}$'s are

$$\lambda'_{B_m(b),0} = f_0(b, q)$$

$$\lambda'_{B_m(b),1} = f_1(b, q) + (b-1)f_2(b, q)$$

$$\lambda'_{B_m(b),2} = f_2(b, q) + 2(b-2)f_3(b, q) + (b-2)(b-3)f_4(b, q)$$

$$\lambda'_{B_m(b),3} = f_3(b, q) + 3(b-3)f_4(b, q) + 3(b-3)(b-4)f_5(b, q) + (b-3)(b-4)(b-5)f_6(b, q)$$

$$\lambda'_{B_m(b),4} = f_4(b, q) + 4(b-4)f_5(b, q) + 6(b-4)(b-5)f_6(b, q) + 4(b-4)(b-5)(b-6)f_7(b, q)$$

$$+(b-4)(b-5)(b-6)(b-7)f_8(b, q) . \quad (4.10)$$

To show that $f_0(b, q) = \lambda_{B_m(b), 0, 1}$ has the largest magnitude for $q \geq b$, we first need $f_i(b, q) > f_{i+1}(b, q) + (b-i-1)f_{i+2}(b, q)$ for $0 \leq i \leq b-2$ and $q \geq b$,

$$\begin{aligned} & f_i(b, q) - f_{i+1}(b, q) - (b-i-1)f_{i+2}(b, q) \\ &= \sum_{s=0}^{b-i} (-1)^s \binom{b-i}{s} (q-i-s)^{(b-i-s)} - \sum_{s=0}^{b-i-1} (-1)^s \binom{b-i-1}{s} (q-i-1-s)^{(b-i-1-s)} \\ & \quad - (b-i-1)f_{i+2}(b, q) \\ &= (-1)^{b-i} + \sum_{s=0}^{b-i-1} (-1)^s (q-i-1-s)^{(b-i-1-s)} \frac{(b-i-1)!}{s!(b-i-s)!} \left[(b-i)(q-i-s) - (b-i-s) \right] \\ & \quad - (b-i-1)f_{i+2}(b, q) \\ &= (-1)^{b-i} + \sum_{s=0}^{b-i-1} (-1)^s (q-i-1-s)^{(b-i-1-s)} \left[\binom{b-i-1}{s} (b-i-1) + \binom{b-i}{s} (q-b) \right] \\ & \quad - (b-i-1)f_{i+2}(b, q) \\ &= (-1)^{b-i} + (-1)^{b-i-1} [b-i-1 + (b-i)(q-b)] + \sum_{s=0}^{b-i-2} (-1)^s (q-i-2-s)^{(b-i-2-s)} \\ & \quad \times \left[(q-i-1-s) \left[\binom{b-i-1}{s} (b-i-1) + \binom{b-i}{s} (q-b) \right] - (b-i-1) \binom{b-i-2}{s} \right] \\ &= (-1)^{b-i} + (-1)^{b-i-1} [b-i-1 + (b-i)(q-b)] + \sum_{s=0}^{b-i-2} (-1)^s (q-i-2-s)^{(b-i-2-s)} \\ & \quad \times \left[(q-b) \left[\binom{b-i-1}{s} (b-i-1) + \binom{b-i}{s} (q-b) + (b-i-1-s) \binom{b-i}{s} \right] \right. \\ & \quad \left. + (b-i-1)(b-i-2) \binom{b-i-2}{s} \right] \\ &\geq 0 \quad \text{for } 0 \leq i \leq b-2 \text{ and } q \geq b , \end{aligned} \quad (4.11)$$

where the equal sign is realized only when $i = b-2$ at $q = b$. Therefore, we have $\lambda'_{B_m(b), 0} > \lambda'_{B_m(b), 1} > \dots > \lambda'_{B_m(b), b/2}$ for even b and $\lambda'_{B_m(b), 0} > \lambda'_{B_m(b), 1} > \dots > \lambda'_{B_m(b), (b-1)/2}$ for odd b . Recall $f_i(b, q) = 0$ if $b < i$. To show $\lambda'_{B_m(b), b/2} > \dots > \lambda'_{B_m(b), b}$ for even b and $\lambda'_{B_m(b), (b+1)/2} > \dots > \lambda'_{B_m(b), b}$ for odd b , we must have the coefficient of $f_b(b, q)$ in $\lambda'_{B_m(b), d}$ is smaller than the sum of the coefficients of $f_{b-2}(b, q)$ and $f_b(b, q)$ in $\lambda'_{B_m(b), d-1}$ as follows,

$$\begin{aligned}
& \binom{d-1}{b-d-1} (b-d+1)^{(b-d-1)} + \binom{d-1}{b-d+1} (b-d+1)^{(b-d+1)} - \binom{d}{b-d} (b-d)^{(b-d)} \\
&= \frac{(d-1)!(b-d+1)!}{(b-d-1)!(2d-b)!2} + \frac{(d-1)!(b-d+1)!}{(b-d+1)!(2d-b-2)!} - \frac{d!(b-d)!}{(b-d)!(2d-b)!} \\
&= \frac{(d-1)!}{(2d-b)!} \frac{(b-d+1)(b-d)}{2} + \frac{(d-1)!}{(2d-b-2)!} - \frac{d!}{(2d-b)!} \\
&= \frac{(d-1)!}{2(2d-b)!} \left[(b-d+1)(b-d) + 2(2d-b)(2d-b-1) - 2d \right] \\
&= \frac{(d-1)!}{2(2d-b)!} (3b^2 - 10bd + 9d^2 + 3b - 7d) , \tag{4.12}
\end{aligned}$$

which is larger than 0 for even b with $b/2 < d \leq b$ and odd b with $(b+1)/2 < d \leq b$ except for $(b, d) = (2, 2)$ and $(4, 3)$ where it is 0. For odd b and $d = (b+1)/2$, there is no $f_b(b, q)$ term in $\lambda'_{B_m(b), (b-1)/2}$, and we need the coefficient of the $f_b(b, q)$ in $\lambda'_{B_m(b), (b+1)/2}$ is smaller than the coefficient of $f_{b-2}(b, q)$ in $\lambda'_{B_m(b), (b-1)/2}$,

$$\begin{aligned}
& \binom{d-1}{b-d-1} (b-d+1)^{(b-d-1)} - \binom{d}{b-d} (b-d)^{(b-d)} \\
&= \frac{(d-1)!(b-d+1)!}{2(b-d-1)!(2d-b)!} - \frac{d!(b-d)!}{(b-d)!(2d-b)!} \\
&= \frac{(d-1)!}{2(2d-b)!} \left[(b-d+1)(b-d) - 2d \right] \\
&= \frac{1}{2} \left(\frac{b-1}{2} \right)! \left[\left(\frac{b+1}{2} \right) \left(\frac{b-1}{2} \right) - (b+1) \right] \\
&= \frac{1}{8} \left(\frac{b-1}{2} \right)! (b-5)(b+1) \geq 0 \quad \text{for } b \geq 5 . \tag{4.13}
\end{aligned}$$

Therefore, for $b \geq 4$, we find $\lambda_{B_m(b), 0, 1}$ is dominant for $q \geq b$, i.e. $q_c(B_m(b)) < b$. $\square$.

We remark that our theorem, in combination with the known results that $q_c(B_m(b)) = b$ for $b = 2, 3$, yields the theorem that

$$q_c(B_m(b)) \leq b \quad \text{for } b \geq 2 . \tag{4.14}$$

We observe that the length between b and $q_c(B_m(b))$ are $-1, 0, 0, 0.3264, 0.4937$, and 0.6764 for the respective values $1 \leq b \leq 6$. These values suggest that this length might well increase monotonically up to 1 as $b \rightarrow \infty$, since the lower bound for $q_c(B_m(b))$ is $b-1$ [62].

We observe the following general form of four $\lambda_{B_m(b), 3, j}$ corresponding to the partition [21],

$$\lambda_{B_m(b),3,j_1} = -f_3(b, q) + 2(b-3)f_4(b, q) - (b-3)(b-4)f_5(b, q) \quad (4.15)$$

$$\lambda_{B_m(b),3,j_2} = -f_3(b, q) + (b-4)f_4(b, q) + (b-4)f_5(b, q) \quad (4.16)$$

$$\lambda_{B_m(b),3,j_3} = -f_3(b, q) + (b-6)f_4(b, q) + 3(b-4)f_5(b, q) \quad (4.17)$$

$$\lambda_{B_m(b),3,j_4} = -f_3(b, q) - 3f_4(b, q) - 3f_5(b, q) . \quad (4.18)$$

These formulae are correct for $3 \leq b \leq 9$, and we conjecture that these are the only $\lambda_{B_m(b),3,j}$ corresponding to the partition [21] for arbitrarily b . For toroidal strips, their coefficients are $c_{B_m(b),3,j} = \frac{1}{6}q(q-2)(q-4)$, and $c'_{B_m(b),3,j_k} = 2(b-1), b(b-3), (b-1)(b-2)$, and $\frac{2}{3}b(b-2)(b-4)$ for $1 \leq k \leq 4$.

From toroidal strips to Klein bottle strips, we observe the transformations from $c'_{B_m(b),d,j}$ to $c'_{\bar{B}_m(b),d,j}$ have certain patterns for $0 \leq d \leq 4$ and $2 \leq b \leq 6$, and summarized them in table XI. Notice that the results are different for even and odd b .

TABLE XI. Relation of $c'_{B_m(b),d,j}$ and $c'_{\bar{B}_m(b),d,j}$

partition	d	$c'_{B_m(b),d,j}$	$c'_{\bar{B}_m(b),d,j}$ for odd b	$c'_{\bar{B}_m(b),d,j}$ for even b
[d]	$d \geq 0$	1	1	1
	$d \geq 1$	$b - 1$	0	-1
	$d \geq 2$	$b(b - 3)/2$	$(b - 3)/2$	$b/2$
	$d \geq 3$	$b(b - 1)(b - 5)/6$	0	$-b/2$
	$d \geq 4$	$b(b - 1)(b - 2)(b - 7)/24$	$(b - 1)(b - 7)/8$	$b(b - 2)/8$
[1^d]	2	$b - 1$	0	-1
	2,3	$(b - 1)(b - 2)/2$	$-(b - 1)/2$	$-(b - 2)/2$
	3,4	$(b - 1)(b - 2)(b - 3)/6$	0	$(b - 2)/2$
	4,5	$(b - 1)(b - 2)(b - 3)(b - 4)/24$	$(b - 1)(b - 3)/8$	$(b - 2)(b - 4)/8$
[21]	3	$2(b - 1)$	0	-2
	3	$b(b - 3)$	$b - 3$	b
	3	$(b - 1)(b - 2)$	$-(b - 1)$	$-(b - 2)$
	3	$2b(b - 2)(b - 4)/3$	2	0
[31]	4	$3(b - 1)$	0	-3
	4	$3b(b - 3)/2$	$3(b - 3)/2$	$3b/2$
	4	$3(b - 1)(b - 2)/2$	$-3(b - 1)/2$	$-3(b - 2)/2$
	4	$b(b - 1)(b - 5)/2$	0	$-3b/2$
	4	$b(b - 2)(b - 4)$	3	0
	4	$3b(b - 1)(b - 3)(b - 6)/8$	$-3(b - 1)(b - 3)/8$	$-3b(b - 6)/8$
[22]	4	$b(b - 3)$	$b - 3$	b
	4	$2b(b - 2)(b - 4)/3$	2	0
	4	$b(b - 1)(b - 4)(b - 5)/6$	$(b - 1)(b - 5)/2$	$b(b - 4)/2$
[211]	4	$3(b - 1)(b - 2)/2$	$-3(b - 1)/2$	$-3(b - 2)/2$
	4	$(b - 1)(b - 2)(b - 3)/2$	0	$3(b - 2)/2$
	4	$b(b - 2)(b - 4)$	3	0
	4	$3b(b - 2)(b - 3)(b - 5)/8$	$-3(b - 3)(b - 5)/8$	$-3b(b - 2)/8$

For the partition $[d]$, if we label $c'_{B_m(b), d \geq 0}$ as the $c'_{B_m(b), d, j} = 1$ in the first row of table XI, and similarly for other $c'_{B_m(b), d \geq d'}$, we conjecture the following forms of the corresponding $c'_{\bar{B}_m(b), d \geq d'}$ for general d' ,

$$c'_{\bar{B}_m(b), d \geq d'} = \begin{cases} \binom{(b-1)/2}{d'/2} - \binom{(b-1)/2}{d'/2-1} & \text{for even } d' \geq 2 \\ 0 & \text{for odd } d' \geq 1 \end{cases} \quad \text{for odd } b, \quad (4.19)$$

and

$$c'_{\bar{B}_m(b), d \geq d'} = \begin{cases} \binom{b/2}{d'/2} & \text{for even } d' \geq 0 \\ -\binom{b/2}{(d'-1)/2} & \text{for odd } d' \geq 1 \end{cases} \quad \text{for even } b. \quad (4.20)$$

For the partition $[1^d]$, if we label $c'_{B_m(b), 2}$ as the $c'_{B_m(b), d, j} = b-1$ in the first row of the $[1^d]$ partition of table XI, and similarly for other $c'_{B_m(b), d'}$, we conjecture the following forms of the corresponding $c'_{\bar{B}_m(b), d'}$ for general d' ,

$$c'_{\bar{B}_m(b), d'} = \begin{cases} (-1)^{(d'-1)/2} \binom{(b-1)/2}{(d'-1)/2} & \text{for odd } d' \geq 3 \\ 0 & \text{for even } d' \geq 2 \end{cases} \quad \text{for odd } b, \quad (4.21)$$

and

$$c'_{\bar{B}_m(b), d'} = \begin{cases} \sum_{j=0}^{(d'-1)/2} (-1)^j \binom{b/2}{j} & \text{for odd } d' \geq 3 \\ \sum_{j=0}^{(d'-2)/2} (-1)^{j-1} \binom{b/2}{j} & \text{for even } d' \geq 2 \end{cases} \quad \text{for even } b, \quad (4.22)$$

If the sum of the coefficients for a specific d is denoted as $C(\bar{B}_m(b), d, q)$, that is,

$$C(\bar{B}_m(b), d, q) = \sum_{j=1}^{N_{\bar{B}_m(b), d, \lambda}} c_{\bar{B}_m(b), d, j}(q) c'_{\bar{B}_m(b), d, j}, \quad (4.23)$$

we have the following conjecture,

$$C(\bar{B}_m(\text{even } b), d, q) = \begin{cases} (-1)^{d/2} \binom{b/2}{d/2} & \text{for even } d \\ 0 & \text{for odd } d \end{cases} \quad (4.24)$$

$$C(\bar{B}_m(\text{odd } b), d, q) = \begin{cases} (-1)^{d/2} \binom{(b-1)/2}{d/2} & \text{for even } d \\ (-1)^{(d-1)/2} (q-1) \binom{(b-1)/2}{(d-1)/2} & \text{for odd } d \end{cases}. \quad (4.25)$$

V. CONCLUSIONS

In this paper we have presented exact zero-temperature partition functions for families of chain graphs composed of repeated complete subgraphs K_b , $b = 5, 6$ with periodic or twisted periodic boundary condition in the longitudinal direction. In the $L_x \rightarrow \infty$ limit, the continuous accumulation set of chromatic zeros $\mathcal{B}$ was determined. We showed the eigenvalues with coefficients of degree $b-1$ are bounded between $(-1)^{b-1}(q-1)$ and $(-1)^{b-1}(q-2b+1)$, and gave the explicit forms of the eigenvalues for partitions of the form [21]. The minimum real q at which $\mathcal{B}$ crosses the real q axis was proven to be $q = 0$, and $q_c(B_m(b)) < b$ for $4 \leq b$.

Acknowledgment: I would like to thank Prof. R. Shrock for helpful discussions.

-
- [1] R. B. Potts, Proc. Camb. Phil. Soc. **48** (1952) 106.
 - [2] F. Y. Wu, Rev. Mod. Phys. **54** (1982) 235.
 - [3] R. C. Read, J. Combin. Theory **4** (1968) 52.
 - [4] R. C. Read and W. T. Tutte, "Chromatic Polynomials", in *Selected Topics in Graph Theory*, 3, eds. L. W. Beineke and R. J. Wilson (Academic Press, New York, 1988.).
 - [5] N. L. Biggs, *Algebraic Graph Theory* (2nd ed., Cambridge Univ. Press, Cambridge, 1993).
 - [6] R. Shrock and S.-H. Tsai, Phys. Rev. **E55** (1997) 5165.
 - [7] N. L. Biggs, LSE report LSE-CDAM-99-08 (1999).
 - [8] S. Beraha, J. Kahane, and N. Weiss, J. Combin. Theory B **27** (1979) 1; *ibid.*, **28** (1980) 52.
 - [9] N. L. Biggs, R. M. Damerell, and D. A. Sands, J. Combin. Theory B **12** (1972) 123.
 - [10] N. L. Biggs and R. Shrock, J. Phys. A (Letts) **32**, L489 (1999).
 - [11] S.-C. Chang, Physica **A296**, 495 (2001).
 - [12] E. H. Lieb, Phys. Rev. **162** (1967) 162.
 - [13] S.-C. Chang and R. Shrock, Phys. Rev. E, **62**, 4650 (2000).
 - [14] S.-C. Chang and R. Shrock, Physica A **296** (2001) 131.
 - [15] N. L. Biggs and G. H. Meredith, J. Combin. Theory B **20** (1976) 5.
 - [16] N. L. Biggs and G. H. Meredith, J. Combin. Theory B **20** (1976) 5.
 - [17] N. L. Biggs, Bull. London Math. Soc. **9** (1977) 54.
 - [18] R. C. Read, in Proc. 3rd Caribbean Conf. on Combin. and Computing (1981).
 - [19] R. C. Read, in Proc. 5th Caribbean Conf. on Combin. and Computing (1988).
 - [20] R. J. Baxter, J. Phys. A **20** (1987) 5241.
 - [21] R. C. Read and G. F. Royle, in *Graph Theory, Combinatorics, and Applications* (Wiley, NY, 1991), vol. 2, p. 1009.
 - [22] R. Shrock and S.-H. Tsai, Phys. Rev. **E55** (1997) 6791.

- [23] R. Shrock and S.-H. Tsai, Phys. Rev. **E56**, 1342, 2733, 3935, 4111 (1997).
- [24] R. Shrock and S.-H. Tsai, Phys. Rev. **E56** (1997) 3935.
- [25] J. Salas and A. Sokal, J. Stat. Phys. **86** (1997) 551.
- [26] M. Roček, R. Shrock, and S.-H. Tsai, Physica **A252** (1998) 505.
- [27] M. Roček, R. Shrock, and S.-H. Tsai, Physica **A259** (1998) 367.
- [28] R. Shrock and S.-H. Tsai, Physica **A259** (1998) 315.
- [29] R. Shrock and S.-H. Tsai, Phys. Rev. **E58** (1998) 4332.
- [30] R. Shrock and S.-H. Tsai, J. Phys. A **31** (1998) 9641; Physica **A265** (1999) 186.
- [31] R. Shrock and S.-H. Tsai, J. Phys. A **32** (1999) 5053.
- [32] R. Shrock and S.-H. Tsai, J. Phys. A Lett. **32** (1999) L195.
- [33] R. Shrock and S.-H. Tsai, Phys. Rev. **E60** (1999) 3512.
- [34] R. Shrock and S.-H. Tsai, Physica A **275** (2000) 429.
- [35] R. Shrock, Phys. Lett. **A261** (1999) 57.
- [36] A. Sokal, Comb. Probab. Comput. **10**, 41 (2001).
- [37] N. L. Biggs, J. Combin. Theory B **82**, 19 (2001); Bull. London Math. Soc., in press.
- [38] R. Shrock, in the *Proceedings of the 1999 British Combinatorial Conference, BCC99*, Discrete Math. **231** (2001) 421.
- [39] H. Kluepfel and R. Shrock, unpublished; H. Kluepfel, Stony Brook thesis “The q -State Potts Model: Partition Functions and Their Zeros in the Complex Temperature and q Planes” (July, 1999).
- [40] R. Shrock, Physica **A281**, 221 (2000).
- [41] S.-C. Chang and R. Shrock, Phys. Rev. **E62**, 4650 (2000).
- [42] R. Shrock, Physica **A283**, 388 (2000).
- [43] S.-C. Chang and R. Shrock, Physica **A286**, 189 (2000).
- [44] S.-C. Chang and R. Shrock, Physica A **290**, 402 (2001).
- [45] S.-C. Chang and R. Shrock, Ann. Phys. **290**, 124 (2001).
- [46] S.-C. Chang and R. Shrock, Phys. Rev. **E62**, 4650 (2000).
- [47] J. Salas and A. Sokal, J. Stat. Phys., **104**, 611 (2001).
- [48] S.-C. Chang and R. Shrock, Physica A **286**, 189 (2000).
- [49] S.-C. Chang and R. Shrock, Physica A **292**, 307 (2001).
- [50] S.-C. Chang and R. Shrock, Physica A **296**, 234 (2001).
- [51] S.-C. Chang and R. Shrock, Physica A **296**, 183 (2001).
- [52] S.-C. Chang and R. Shrock, Int. J. Mod. Phys. B **15**, 443 (2001).
- [53] J. L. Jacobsen and J. Salas, J. Stat. Phys. **104**, 703 (2001).
- [54] J. Salas and R. Shrock, Phys. Rev. E **64**, 011111 (2001).
- [55] S.-C. Chang and R. Shrock, Physica A, **301**, 301 (2001).
- [56] S.-C. Chang and R. Shrock, Phys. Rev. E, in press; cond-mat/0107012.
- [57] J. Salas, S.-C. Chang and R. Shrock, J. Stat. Phys., in press (cond-mat/0108144).
- [58] N. L. Biggs, “Chromatic polynomials and representations of the symmetric group”, AGT workshop, Edinburgh (July 2001), LSE report LSE-CDAM-01-02 (2001).
- [59] N. L. Biggs, M. Klin, and P. Reinfeld, “Algebraic methods for chromatic polynomials”, AGT workshop, Edinburgh (July, 2001), LSE report LSE-CDAM-01-06 (2001).

- [60] N. L. Biggs, LSE report LSE-CDAM-00-04 (May 2000).
- [61] N. L. Biggs and P. Reinfeld, LSE report LSE-CDAM-00-07 (June 2000),
<http://www.cdam.lse.ac.uk/Reports/> .
- [62] N. L. Biggs, S.-C. Chang, P. Reinfeld, and R. Shrock, in preparation.